





[Name of Document] Application for Patent
[Date of Submission] Year Month Day
Heisei 14 (2002) 6 17
[Ref. No.] P7194GN
[Addressed to] Commissioner, Patent Office
[IPC] C12Q 1/68
C07H 21/04
[Inventor]
[Address] c/o Banyu Pharmaceutical Co., Ltd. Tukuba Research Institute, 3, Okubo, Tsukuba-shi, Ibaraki JAPAN
[Name] KOTANI, Hidehito
[Inventor]
[Address] c/o Banyu Pharmaceutical Co., Ltd. Tukuba Research Institute, 3, Okubo, Tsukuba-shi, Ibaraki JAPAN
[Name] MIZUARAI, Shinji
[Applicant]
[Identification No.] 000005072
[Name] Banyu Pharmaceutical Co., Ltd.
[Patent Attorney]
[Identification No.] 100080816
[Name] KATO, Asamichi, Patent Attorney
[Fee]
[Amount] Yen 21,000
[List of Documents]
[Name of document] Description 1
[Name of document] Drawings 1
[Name of document] Abstract 1

[Fig. 2]



[Name of Document] Drawings
[Fig. 1]



[Name of Document]

ABSTRACT

[Abstract]

[Problems]

To provide polymorphisms of ABCG2 polypeptide and polynucleotide coding therefor, which is related to the intracellular accumulation of indolocarbazole compounds, as well as methods for detecting the polymorphisms.

[Means for Solution]

A sample is collected from mammals, and a polymorphism of the nucleotide sequence of *ABCG2* gene or a polymorphism of the amino acid sequence of ABCG2 polypeptide is determined. In a preferred embodiment of the present invention, the polymorphism of the nucleotide sequence is one or more of single nucleotide polymorphisms at positions selected from the group consisting of 34, 376 and 421 of SEQ ID NO:1, and the polymorphism of the amino acid sequence is one or more of amino acid polymorphisms at positions consisting of 12, 126, and 141 of SEQ ID NO:2.

[Selected Drawing]

Fig. 1